

κ -DEFORMED OSCILLATORS, THE CHOICE OF STAR PRODUCT AND FREE κ -DEFORMED QUANTUM FIELDS

Marcin Daszkiewicz, Jerzy Lukierski, Mariusz Woronowicz

Institute of Theoretical Physics

University of Wrocław pl. Maxa Born'a 9, 50-206 Wrocław, Poland

e-mail: marcin,lukier,woronow@ift.uni.wroc.pl

Abstract

The aim of this paper is to study the general framework providing various κ -deformations of field oscillators and consider the commutator function of the corresponding κ -deformed free fields. In order to obtain free κ -deformed quantum fields (with c -number commutators) we proposed earlier a particular model of κ -deformed oscillator algebra [1] and the modification of κ -star product [2], implementing in the product of two quantum fields the change of standard κ -deformed mass-shell conditions. We recall here that other different models of κ -deformed oscillators recently introduced in [3]-[5] are defined on standard κ -deformed mass-shell. In this paper we consider the most general κ -deformed field oscillators, parametrized by set of arbitrary functions in three-momentum space. Firstly, we study the fields with the κ -deformed oscillators defined on standard κ -deformed mass-shell, and argue that for any such a choice of κ -deformed field oscillators algebra we do not obtain the free quantum κ -deformed fields with the c -number commutators. Further, we study κ -deformed quantum fields with the modified κ -star product and derive large class of κ -oscillators defined on suitably modified κ -deformed mass-shell. We obtain large class of κ -deformed statistics depending on six arbitrary functions which all provide the c -number field commutator functions. This general class of κ -oscillators can be described by the composition of suitably defined κ -multiplications and the κ -deformation of the flip operator.

1 Introduction

The standard relativistic local quantum fields on Minkowski space provide a basic tool for the description of fundamental interactions. If we include into consideration the quantum gravity effects such classical description breaks down at Planck distances ($\approx 10^{-33}$ cm) and it is quite plausible that space-time becomes noncommutative (see e.g. [6], [7]). The κ -deformation of Minkowski space [8]-[10] and corresponding quantum κ -deformation of relativistic symmetries (see e.g. [11], [12], [9]) provides a fundamental mass scale and possible tools for the description of Planckian quantum-gravitational regime.

At present an important task is the construction of κ -deformed quantum field theory on κ -deformed Minkowski space. In contrast with simpler, recently studied case of θ -deformed symmetries (with constant commutator $[\hat{x}_\mu, \hat{x}_\nu] = \theta_{\mu\nu}$; see e.g. [13], [14]), the κ -deformation of relativistic symmetries is not described by a twist factor¹, and for the κ -Poincare algebra the universal R-matrix is not known. We can therefore apply mainly the technique of star product (for application to κ -deformation see e.g. [17]-[22]) as representing the κ -deformation of space-time.

a) *Summary of previous results* [1], [2], [25]. Main aim of our scheme presented earlier in [1], [2], [25] was the construction of free quantum κ -deformed fields characterized by c -number commutator function². It should be stressed that for such free κ -deformed quantum fields all other properties known from the standard free quantum field theory as the notion of locality, microcausality, the structure of Fock space, or kinematic independence of field excitations defining the multiparticle states are modified. The c -number commutator of quantum κ -deformed fields is obtained by the interplay of the two following sources of noncommutativity: quantum nature of space-time and specific κ -deformation of the field oscillators algebra:

i) *Noncommutativity of space-time*. We replace the standard quantum field arguments x_μ by κ -Minkowski noncommutative coordinates $\hat{x}_\mu$

$$[\hat{x}_0, \hat{x}_i] = \frac{i}{\kappa} \hat{x}_i, \quad [\hat{x}_i, \hat{x}_j] = 0. \quad (1)$$

One can introduce the Fourier expansion of κ -deformed free quantum fields

$$\hat{\varphi}(\hat{x}) = \frac{1}{(2\pi)^{\frac{3}{2}}} \int d^4p \hat{A}(p_0, \vec{p}) \delta(C_2^\kappa(\vec{p}, p_0) - M^2) :e^{ip_\mu \hat{x}^\mu}:, \quad (2)$$

where the symmetrized κ -deformed plane wave looks as follows (see e.g. [26])

$$:e^{ip_\mu \hat{x}^\mu}: = e^{\frac{i}{2}p_0 \hat{x}_0} e^{ip_i \hat{x}_i} e^{\frac{i}{2}p_0 \hat{x}_0}. \quad (3)$$

The operators $\hat{A}(p_0, \vec{p})$ describe the quantized field oscillators and $C_2^\kappa(\vec{p}, p_0)$ represents the κ -deformed mass square Casimir

$$C_2^\kappa(\vec{p}, p_0) = \left(2\kappa \sinh\left(\frac{p_0}{2\kappa}\right)\right)^2 - \vec{p}^2, \quad (4)$$

¹We restrict here ourselves to twists $T \in \mathcal{U}(P_4) \otimes \mathcal{U}(P_4)$ where $\mathcal{U}(P_4)$ describes the enveloping $D = 4$ Poincare algebra. For twists not satisfying this condition see [15], [16].

²We extend to noncommutative quantum fields the known definition of (generalized) free fields as described by a c -number commutator (see e.g. [23], [24]).

defining the κ -deformed mass-shell condition

$$C_2^\kappa(\vec{p}, p_0) - M^2 = 0 , \quad (5)$$

which implies the energy-momentum dispersion relation

$$p_0 = \pm \omega_\kappa(\vec{p}) , \quad (6)$$

where

$$\omega_\kappa(\vec{p}) = 2\kappa \operatorname{arcsinh} \left(\frac{\omega(\vec{p})}{2\kappa} \right) \quad ; \quad \omega(\vec{p}) = \sqrt{\vec{p}^2 + M^2} . \quad (7)$$

For the discussion of κ -deformed free fields we introduce the operator-valued Weyl homomorphism

$$\hat{\varphi}(\hat{x}) \leftrightarrow \hat{\varphi}(x) \quad , \quad \hat{\varphi}(\hat{y}) \leftrightarrow \hat{\varphi}(y) , \quad (8)$$

where the relations (1) describing the pair of noncommutative κ -Minkowski points $\hat{x}$, $\hat{y}$ is supplemented by the following additional crossrelations³

$$[\hat{x}_0, \hat{y}_i] = \frac{i}{\kappa} \hat{y}_i \quad , \quad [\hat{y}_0, \hat{x}_i] = \frac{i}{\kappa} \hat{x}_i . \quad (9)$$

The corresponding κ -star multiplication prescription which represents the noncommutative space-time structure (1), (9) looks as follows

$$\begin{aligned} \hat{\varphi}(\hat{x}) \cdot \hat{\varphi}(\hat{y}) \leftrightarrow \hat{\varphi}(x) \star_\kappa \hat{\varphi}(y) &= \frac{1}{(2\pi)^3} \int d^4 p \int d^4 q \, e^{i(p_0 x_0 + q_0 y_0) + (p_i e^{\frac{q_0}{2\kappa}} x^i + q_i e^{-\frac{p_0}{2\kappa}} y^i)} \\ &\cdot \hat{A}(p_0, \vec{p}) \hat{A}(q_0, \vec{q}) \delta(C_\kappa^2(\vec{p}, p_0) - M^2) \delta(C_\kappa^2(\vec{q}, q_0) - M^2) . \end{aligned} \quad (10)$$

The formula (10) follows from the κ -star products of exponentials (3)

$$\colon e^{ip_\mu \hat{x}^\mu} \colon \cdot \colon e^{iq_\mu \hat{y}^\mu} \colon \leftrightarrow e^{ip_\mu x^\mu} \star_\kappa e^{iq_\mu y^\mu} = e^{i(p_0 x^0 + q_0 y^0) + (p_i e^{\frac{q_0}{2\kappa}} x^i + q_i e^{-\frac{p_0}{2\kappa}} y^i)} , \quad (11)$$

where rhs of (11) is determined by the fourmomentum coproduct

$$\Delta(P_0) = P_0 \otimes 1 + 1 \otimes P_0 \quad , \quad \Delta(P_i) = P_i \otimes e^{\frac{P_0}{2\kappa}} + e^{-\frac{P_0}{2\kappa}} \otimes P_i . \quad (12)$$

The field oscillators in the product $\hat{\varphi}(x) \star_\kappa \hat{\varphi}(y)$ (see (10)) carry the fourmomenta satisfying the standard κ -deformed mass-shell condition (5). We recall that in the models of κ -statistics presented in [3]-[5] there is imposed such a standard κ -deformed mass-shell condition.

In our papers [2], [25] we modified the standard κ -star product (10) by the following additional deformation of the product of mass-shell deltas

$$\delta(C_\kappa^2(\vec{p}, p_0) - M^2) \delta(C_\kappa^2(\vec{q}, q_0) - M^2) \Rightarrow \delta(C_\kappa^2(\vec{p} e^{\frac{q_0}{2\kappa}}, p_0) - M^2) \delta(C_\kappa^2(\vec{q} e^{-\frac{p_0}{2\kappa}}, q_0) - M^2) . \quad (13)$$

³The relation (9) permits to perform in consistency with (1) the limit $\hat{x}_\mu \rightarrow \hat{y}_\mu$. Other option for constructing the κ -deformed field theory is to assume that $[\hat{x}_\mu, \hat{y}_\nu] = 0$ (see e.g. [27]).

Such modification can be also introduced as the change of standard multiplication of the fields on noncommutative space-time

$$\hat{\varphi}(\hat{x}) \cdot \hat{\varphi}(\hat{y}) \rightarrow \hat{\varphi}(\hat{x}) \cdot_{\kappa} \hat{\varphi}(\hat{y}) = \frac{1}{(2\pi)^3} \int d^4 p \int d^4 q \hat{A}(p_0, \vec{p}) \hat{A}(q_0, \vec{q}) e^{i(p\hat{x} + q\hat{y})} \cdot \delta(C_{\kappa}^2(\vec{p}e^{\frac{q_0}{2\kappa}}, p_0) - M^2) \delta(C_{\kappa}^2(\vec{q}e^{-\frac{p_0}{2\kappa}}, q_0) - M^2), \quad (14)$$

which is homomorphic to the following modified κ -star product of two κ -deformed free fields on commuting Minkowski space

$$\hat{\varphi}(x) \hat{\star}_{\kappa} \hat{\varphi}(y) = \frac{1}{(2\pi)^3} \int d^4 p \int d^4 q \hat{A}(p_0, \vec{p}) \hat{A}(q_0, \vec{q}) e^{i(p_0 x_0 + q_0 y_0) + (p_i e^{\frac{q_0}{2\kappa}} x^i + q_i e^{-\frac{p_0}{2\kappa}} y^i)} \cdot \delta(C_{\kappa}^2(\vec{p}e^{\frac{q_0}{2\kappa}}, p_0) - M^2) \delta(C_{\kappa}^2(\vec{q}e^{-\frac{p_0}{2\kappa}}, q_0) - M^2), \quad (15)$$

ii) *κ -deformation of field oscillators algebra.* We see from (15) that the field oscillators in the product $\hat{\varphi}(x) \hat{\star}_{\kappa} \hat{\varphi}(y)$ are put on modified mass-shells (see (13)). The unconventional feature of such an approach is the use of quantized field oscillators $\hat{A}(p_0, \vec{p})$ extended to the values of $p = (p_0, \vec{p})$ which do not satisfy the κ -deformed mass-shell condition (4)

$$\hat{A}(p_0, \vec{p})|_{C_{\kappa}^2(\vec{p}, p_0)=M^2} \Rightarrow \hat{A}(p_0, \vec{p})|_{C_{\kappa}^2(\vec{p}, p_0) \neq M^2}. \quad (16)$$

The pair of modified mass-shell conditions following from (13) is

$$C_{\kappa}^2(\vec{p}e^{\frac{q_0}{2\kappa}}, p_0) - M^2 = 0 \quad , \quad C_{\kappa}^2(\vec{q}e^{-\frac{p_0}{2\kappa}}, q_0) - M^2 = 0. \quad (17)$$

Such a modification can be interpreted by the coproduct relation (12), where $(p_0, \vec{p}e^{q_0/2\kappa})$ and $(q_0, \vec{q}e^{-p_0/2\kappa})$ correspond to first and second term in the fourmomentum addition formula for the two-particle state

$$P_0 \triangleright \hat{A}(p) \hat{A}(q) = (p_0 + q_0) \hat{A}(p) \hat{A}(q), \quad (18)$$

$$P_i \triangleright \hat{A}(p) \hat{A}(q) := m \circ [\Delta(P_i) \hat{A}(p) \otimes \hat{A}(q)] = [p_i e^{q_0/2\kappa} + q_i e^{-p_0/2\kappa}] \hat{A}(p) \hat{A}(q). \quad (19)$$

where $m \circ (f \otimes g) = fg$. The energy values p_0, q_0 satisfying (17) are equivalently described as the solutions $p_0^{(\epsilon, \epsilon')}(\vec{p}, \vec{q})$, $q_0^{(\epsilon, \epsilon')}(\vec{p}, \vec{q})$ ($\epsilon = \pm 1$, $\epsilon' = \pm 1$) of the following two coupled equations

$$p_0^{(\epsilon, \epsilon')} = \epsilon \omega_{\kappa}(\vec{p}e^{q_0^{(\epsilon, \epsilon')}/2\kappa}) \quad , \quad q_0^{(\epsilon, \epsilon')} = \epsilon' \omega_{\kappa}(\vec{q}e^{-p_0^{(\epsilon, \epsilon')}/2\kappa}). \quad (20)$$

We see from (20) that for κ -deformed two-particle states the energy of the first particle depends also on the three-momenta of the energy, i.e. the modification (13) of the mass-shell condition couples both constituents of the two-particle state.

For particular choice of κ -deformed oscillator algebra we did show in [2], [25] that using the modified κ -star product (15) one gets the c -number field commutator. One obtains

$$[\hat{\varphi}(x), \hat{\varphi}(y)]_{\hat{\star}_{\kappa}} = i \Delta_{\kappa}(x - y; M^2), \quad (21)$$

where

$$\Delta_{\kappa}(x; M^2) = \frac{i}{(2\pi)^3} \int d^4 p \epsilon(p_0) \delta((2\kappa \sinh(p_0/2\kappa))^2 - \vec{p}^2 - M^2) e^{ip_{\mu} x^{\mu}}, \quad (22)$$

is the κ -deformed Pauli-Jordan commutator function proposed firstly in [12] after using somewhat naive arguments.

b) *Plan of the paper and the resume of results.* In our earlier papers [1], [2], [25] we studied a definite model of κ -deformed oscillators, without any free parameter (except the deformation parameter κ). In this paper we consider the most general set of binary κ -deformed oscillator algebras consistent with κ -deformed addition law of the fourmomenta.

Firstly, in Sect. 2 we shall use κ -star product (10) and the oscillators $\hat{A}(p_0, \vec{p})$ which lie on standard κ -deformed mass-shells. We shall show that in such a case for any possible choice of binary oscillator algebra it is not possible to obtain the c -number value of the field commutator. In place of the formula (21) one gets the q -number field commutator bilinear in the κ -deformed oscillators. In particular we shall discuss briefly two recent proposals of κ -deformed oscillator algebras ([3], [4]) which fall into such a category.

In Sect. 3 we shall consider the binary multiplication of κ -deformed free fields using the modified κ -deformed $\star$ -product. In such a case by performing the general transformation in two-particle fourmomentum space we arrive at large class of κ -deformed oscillator algebras depending on six arbitrary functions of two-particle fourmomenta, which all lead to the c -number field commutator. All such κ -oscillators are characterized by modified energy-momentum dispersion relation generalizing the relations (17) or (20) and can be classified by various forms of the addition law for the threemomenta of κ -deformed two-particle state. The particular choice proposed in [1], [2] was described by the Abelian addition law

$$\vec{p}_{1+2} = \vec{p} + \vec{q}, \quad (23)$$

while the choices from [3], [4] (see also [16]) were characterized by the non-Abelian addition formula⁴

$$\vec{p}_{1+2} = \vec{p} \dot{+} \vec{q} = \vec{p} e^{\frac{q_0}{2\kappa}} + \vec{q} e^{-\frac{p_0}{2\kappa}}. \quad (24)$$

The κ -deformed oscillator algebra with the threemomentum addition law (24) was constructed by the use of suitable deformation of the flip operator. In Sect. 4 we describe the general oscillator algebras (61) (see Sect. 3) as the composition of the most general κ -deformed multiplication and the κ -deformed flip operation.

The κ -deformed field oscillators determine the structure of corresponding κ -deformed multiparticle states⁵. We shall only mention here that the n -particle sector of κ -deformed Fock space is represented by suitably constructed non-factorizable clusters. We interpret such structure of κ -deformed Fock spaces as a result of the geometric interactions implied by the Lie-algebraic noncommutativity of κ -deformed space-time. It would be interesting to understand how such features can be linked with quantum gravity framework.

⁴We recall that we use the κ -Poincare algebra which leads to the standard (see [11]) coproduct (12) for the threemomenta.

⁵The discussion of multiparticle states for $n > 2$ has been recently considered in [28].

2 Standard κ -star product and the κ -deformed quantum fields

The aim of this section is to study the commutator of the fields (2) with the κ -star multiplication rule (10)

$$\begin{aligned} [\hat{\varphi}(x), \hat{\varphi}(y)]_{\star_\kappa} &= \\ &= \frac{1}{(2\pi)^3} \int d^4p d^4q \hat{A}(p) \hat{A}(q) e^{ipx} \star_\kappa e^{iqy} \delta(C_\kappa^2(p) - M^2) \delta(C_\kappa^2(q) - M^2) \\ &\quad - \frac{1}{(2\pi)^3} \int d^4p' d^4q' \hat{A}(q') \hat{A}(p') e^{iq'y} \star_\kappa e^{ip'x} \delta(C_\kappa^2(p') - M^2) \delta(C_\kappa^2(q') - M^2) . \end{aligned} \quad (25)$$

We see from (25) that due to the presence of respective Dirac deltas the field oscillators remain on κ -deformed mass-shell (4). We shall look for the generalized binary relations of κ -deformed oscillators consistent with the κ -deformed fourmomentum addition law. We shall show that for any choice of these binary relations determining the choice of κ -statistics, the commutator (25) is an operator bilinear in the field oscillators. We use in this paper the definition of (generalized) free field as characterized by the c -number commutator function (see e.g. [23], [24]). We obtain therefore in this section the result that using standard on-shell κ -oscillators $\hat{A}(p_0, \vec{p})$ we can not obtain the free κ -deformed quantum fields.

Our demonstration of the operator nature of the commutator (25) follows from the impossibility of the factorization under momenta integrals of any binary relations for the field oscillators $\hat{A}(p_0, \vec{p})$. For studying possible factorization we shall perform the following general O(3)-covariant change of variables separately in the first term on rhs of (25) ($\tilde{f} = \tilde{f}(p, q)$ etc)

$$\vec{p} \rightarrow \vec{\mathcal{P}}(p, q) = \vec{p}\tilde{f} + \vec{q}\tilde{g} \quad , \quad \vec{q} \rightarrow \vec{\mathcal{Q}}(p, q) = \vec{p}\tilde{k} + \vec{q}\tilde{l} \quad , \quad (26)$$

$$p_0 \rightarrow \mathcal{P}_0(p, q) \quad , \quad q_0 \rightarrow \mathcal{Q}_0(p, q) \quad , \quad (27)$$

with the following inverse formulae ($f = f(\mathcal{P}, \mathcal{Q})$ etc)

$$\vec{\mathcal{P}} \rightarrow \vec{p}(\mathcal{P}, \mathcal{Q}) = \vec{\mathcal{P}}f + \vec{\mathcal{Q}}g \quad , \quad \vec{\mathcal{Q}} \rightarrow \vec{q}(\mathcal{P}, \mathcal{Q}) = \vec{\mathcal{P}}k + \vec{\mathcal{Q}}l \quad , \quad (28)$$

$$\mathcal{P}_0 \rightarrow p_0(\mathcal{P}, \mathcal{Q}) \quad , \quad \mathcal{Q}_0 \rightarrow q_0(\mathcal{P}, \mathcal{Q}) \quad , \quad (29)$$

and in the second term on rhs of (25)

$$\vec{p}' \rightarrow \vec{\mathcal{P}}' = \vec{p}'\tilde{f}' + \vec{q}'\tilde{g}' \quad , \quad \vec{q}' \rightarrow \vec{\mathcal{Q}}' = \vec{p}'\tilde{k}' + \vec{q}'\tilde{l}' \quad , \quad (30)$$

$$p'_0 \rightarrow \mathcal{P}'_0 \quad , \quad q'_0 \rightarrow \mathcal{Q}'_0 \quad , \quad (31)$$

with the inverse formulae

$$\vec{\mathcal{P}}' \rightarrow \vec{p}'(\mathcal{P}', \mathcal{Q}') = \vec{\mathcal{P}}'f' + \vec{\mathcal{Q}}'g' \quad , \quad \vec{\mathcal{Q}}' \rightarrow \vec{q}'(\mathcal{P}', \mathcal{Q}') = \vec{\mathcal{P}}'k' + \vec{\mathcal{Q}}'l' \quad , \quad (32)$$

$$\mathcal{P}_0 \rightarrow p'_0(\mathcal{P}', \mathcal{Q}') \quad , \quad \mathcal{Q}_0 \rightarrow q'_0(\mathcal{P}', \mathcal{Q}') \quad . \quad (33)$$

We assume that f, g, h, k, p_0, q_0 in (28) and $f', g', h', k', p'_0, q'_0$ in (32) are respectively arbitrary $O(3)$ -invariant functions of the fourmomenta $\mathcal{P} = (\vec{\mathcal{P}}, \mathcal{P}_0)$, $\mathcal{Q} = (\vec{\mathcal{Q}}, \mathcal{Q}_0)$ and $\mathcal{P}' = (\vec{\mathcal{P}}', \mathcal{P}'_0)$, $\mathcal{Q}' = (\vec{\mathcal{Q}}', \mathcal{Q}'_0)$ ⁶.

We shall look for such a choice of arbitrary functions in the formulae (28),(29) and (32), (33) which leads to the equality

$$e^{ip(\mathcal{P}, \mathcal{Q})x} \star_\kappa e^{iq(\mathcal{P}, \mathcal{Q})y} = e^{iq'(\mathcal{P}, \mathcal{Q})y} \star_\kappa e^{ip'(\mathcal{P}, \mathcal{Q})x} , \quad (34)$$

with κ -deformed mass-shell conditions taken into account. The formula (34) describes a necessary condition which permits to factorize in the commutator (25) the oscillator algebra and to derive the c -number commutator function. We should observe that the change of variables (26)-(32) modifies as well the explicit form of the product of mass-shell deltas in (25). We obtain in first term of rhs of (25)

$$\begin{aligned} \delta(C_\kappa^2(p) - M^2) \cdot \delta(C_\kappa^2(q) - M^2) &\rightarrow \\ &\rightarrow \delta(C_\kappa^2(p(\mathcal{P}, \mathcal{Q})) - M^2) \cdot \delta(C_\kappa^2(q(\mathcal{P}, \mathcal{Q})) - M^2) , \end{aligned} \quad (35)$$

and in the second term we get

$$\begin{aligned} \delta(C_\kappa^2(p') - M^2) \cdot \delta(C_\kappa^2(q') - M^2) &\rightarrow \\ &\rightarrow \delta(C_\kappa^2(p'(\mathcal{P}, \mathcal{Q})) - M^2) \cdot \delta(C_\kappa^2(q'(\mathcal{P}, \mathcal{Q})) - M^2) . \end{aligned} \quad (36)$$

One obtains⁷

$$\begin{aligned} [\hat{\varphi}(x), \hat{\varphi}(y)]_{\star_\kappa} &= \frac{1}{(2\pi)^3} \int d^4\mathcal{P} d^4\mathcal{Q} J \left(\begin{smallmatrix} p, q \\ \mathcal{P}, \mathcal{Q} \end{smallmatrix} \right) \hat{A}(p(\mathcal{P}, \mathcal{Q})) \hat{A}(q(\mathcal{P}, \mathcal{Q})) \\ &\quad \cdot e^{ip(\mathcal{P}, \mathcal{Q})x} \star_\kappa e^{iq(\mathcal{P}, \mathcal{Q})y} \\ &\quad \cdot \delta(C_\kappa^2(p(\mathcal{P}, \mathcal{Q})) - M^2) \cdot \delta(C_\kappa^2(q(\mathcal{P}, \mathcal{Q})) - M^2) \\ &\quad - \frac{1}{(2\pi)^3} \int d^4\mathcal{P} d^4\mathcal{Q} J \left(\begin{smallmatrix} p', q' \\ \mathcal{P}, \mathcal{Q} \end{smallmatrix} \right) \hat{A}(q'(\mathcal{P}, \mathcal{Q})) \hat{A}(p'(\mathcal{P}, \mathcal{Q})) \\ &\quad \cdot e^{iq'(\mathcal{P}, \mathcal{Q})y} \star_\kappa e^{ip'(\mathcal{P}, \mathcal{Q})x} \\ &\quad \cdot \delta(C_\kappa^2(p'(\mathcal{P}, \mathcal{Q})) - M^2) \cdot \delta(C_\kappa^2(q'(\mathcal{P}, \mathcal{Q})) - M^2) . \end{aligned} \quad (37)$$

Let us denote by $\mathcal{P}_0 = \pi(\vec{\mathcal{P}}, \vec{\mathcal{Q}})$, $\mathcal{Q}_0 = \rho(\vec{\mathcal{P}}, \vec{\mathcal{Q}})$ and $\mathcal{P}'_0 = \pi'(\vec{\mathcal{P}}, \vec{\mathcal{Q}})$, $\mathcal{Q}'_0 = \rho'(\vec{\mathcal{P}}, \vec{\mathcal{Q}})$ the solutions of the following two pairs of the deformed mass-shell conditions

$$C_\kappa^2(p(\mathcal{P}, \mathcal{Q})) - M^2 = 0 \quad , \quad C_\kappa^2(q(\mathcal{P}, \mathcal{Q})) - M^2 = 0 , \quad (38)$$

$$C_\kappa^2(p'(\mathcal{P}, \mathcal{Q})) - M^2 = 0 \quad , \quad C_\kappa^2(q'(\mathcal{P}, \mathcal{Q})) - M^2 = 0 . \quad (39)$$

⁶We recall that the classical $SO(3)$ Hopf algebra is a sub-Hopf algebra of κ -deformed Poincare Hopf algebra.

⁷Because in transformed formula (25) we integrate over the variables $\mathcal{P}'$, $\mathcal{Q}'$, further we shall denote them similarly as in first term of the commutator by $\mathcal{P}$ and $\mathcal{Q}$.

We shall consider firstly the validity of restricted relation (34) obtained by putting $x_0 = y_0 = 0$. It is easy to check that one gets the equality of deformed threemomenta exponentials

$$\begin{aligned}
& e^{ip(\mathcal{P}, \mathcal{Q})x} \star_{\kappa} e^{iq(\mathcal{P}, \mathcal{Q})y} \Big|_{\mathcal{P}_0=\pi, \mathcal{Q}_0=\rho, (x_0=y_0=0)} \\
&= \exp \left[i(\vec{p}(\mathcal{P}, \mathcal{Q}) e^{\frac{q_0(\mathcal{P}, \mathcal{Q})}{2\kappa}} \vec{x} + \vec{q}(\mathcal{P}, \mathcal{Q}) e^{-\frac{p_0(\mathcal{P}, \mathcal{Q})}{2\kappa}} \vec{y}) \right] \Big|_{\mathcal{P}_0=\pi, \mathcal{Q}_0=\rho} \\
&= \exp \left[i(\vec{q}'(\mathcal{P}, \mathcal{Q}) e^{\frac{p'_0(\mathcal{P}, \mathcal{Q})}{2\kappa}} \vec{y} + \vec{p}'(\mathcal{P}, \mathcal{Q}) e^{-\frac{q'_0(\mathcal{P}, \mathcal{Q})}{2\kappa}} \vec{x}) \right] \Big|_{\mathcal{P}_0=\pi', \mathcal{Q}_0=\rho'} \\
&= e^{iq'(\mathcal{P}, \mathcal{Q})y} \star_{\kappa} e^{ip'(\mathcal{P}, \mathcal{Q})x} \Big|_{\mathcal{P}'_0=\pi', \mathcal{Q}'_0=\rho', (x_0=y_0=0)} ,
\end{aligned} \tag{40}$$

if the arbitrary functions introduced in the relations (28),(29) and (32),(33) satisfy the relations

$$\begin{aligned}
f(\mathcal{P}, \mathcal{Q}) e^{p_0(\mathcal{P}, \mathcal{Q})/2\kappa} &= f'(\mathcal{P}, \mathcal{Q}) e^{-p'_0(\mathcal{P}, \mathcal{Q})/2\kappa} , \quad g(\mathcal{P}, \mathcal{Q}) e^{p_0(\mathcal{P}, \mathcal{Q})/2\kappa} = g'(\mathcal{P}, \mathcal{Q}) e^{-p'_0(\mathcal{P}, \mathcal{Q})/2\kappa} , \\
k(\mathcal{P}, \mathcal{Q}) e^{-q_0(\mathcal{P}, \mathcal{Q})/2\kappa} &= k'(\mathcal{P}, \mathcal{Q}) e^{q'_0(\mathcal{P}, \mathcal{Q})/2\kappa} , \quad l(\mathcal{P}, \mathcal{Q}) e^{-q_0(\mathcal{P}, \mathcal{Q})/2\kappa} = l'(\mathcal{P}, \mathcal{Q}) e^{q'_0(\mathcal{P}, \mathcal{Q})/2\kappa} ,
\end{aligned} \tag{41}$$

and $\mathcal{P}_0 = \pi, \mathcal{Q}_0 = \rho, \mathcal{P}_0 = \pi', \mathcal{Q}_0 = \rho'$ satisfy the mass-shell conditions (38),(39).

If we assume the relations (41), then the formula (34) reduces to the equality of time exponentials

$$e^{i[x_0 p_0(\mathcal{P}, \mathcal{Q}) + y_0 q_0(\mathcal{P}, \mathcal{Q})]} \Big|_{\mathcal{P}_0=\pi, \mathcal{Q}_0=\rho} = e^{i[x_0 p'_0(\mathcal{P}, \mathcal{Q}) + y_0 q'_0(\mathcal{P}, \mathcal{Q})]} \Big|_{\mathcal{P}_0=\pi', \mathcal{Q}_0=\rho'} . \tag{42}$$

The relation (42) is satisfied for any values of x_0, y_0 only if

$$p_0(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = p'_0(\vec{\mathcal{P}}, \pi'; \vec{\mathcal{Q}}, \rho') \quad , \quad q_0(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = q'_0(\vec{\mathcal{P}}, \pi'; \vec{\mathcal{Q}}, \rho') . \tag{43}$$

In order to derive the restrictions following from (43) one can rewrite the relations (38),(39) in the form of the following identities ($\epsilon = \pm 1$ etc)

$$p_0(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = \epsilon \omega_{\kappa}(\vec{p}^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho)) \quad , \quad q_0(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = \eta \omega_{\kappa}(\vec{q}^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho)) , \tag{44}$$

$$p'_0(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = \epsilon' \omega_{\kappa}(\vec{p}'^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho)) \quad , \quad q'_0(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = \eta' \omega_{\kappa}(\vec{q}'^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho)) , \tag{45}$$

where $\omega_{\kappa}(\vec{p})$ is defined by relation (7). We see from (44),(45) that the relation (43) can be written as follows⁸

$$\omega_{\kappa}(\vec{p}^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho)) = \omega_{\kappa}(\vec{p}'^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho)) , \tag{46}$$

$$\omega_{\kappa}(\vec{q}^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho)) = \omega_{\kappa}(\vec{q}'^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho)) , \tag{47}$$

which implies that

$$\vec{p}^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = \vec{p}'^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) \quad , \quad \vec{q}^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = \vec{q}'^{\ 2}(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) . \tag{48}$$

Inserting in first formula of (48) the relations (28),(29) and (32),(33) one gets from (46) the condition

$$f^2 \vec{\mathcal{P}}^2 + 2fg \vec{\mathcal{P}} \vec{\mathcal{Q}} + g^2 \vec{\mathcal{Q}}^2 = f'^2 \vec{\mathcal{P}}^2 + 2f'g' \vec{\mathcal{P}} \vec{\mathcal{Q}} + g'^2 \vec{\mathcal{Q}}^2 , \tag{49}$$

⁸Because $\omega_{\kappa} \geq 0$ the relations (43) require that $\epsilon = \epsilon'$ and $\eta = \eta'$.

and analogous relation obtained by the replacements $\vec{p} \rightarrow \vec{p}'$ and $\vec{q} \rightarrow \vec{q}'$ in (47). The insertion of (41) into (49) leads to

$$(1 - e^{(p_0+p'_0)/2\kappa})\vec{p}^2(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = 0. \quad (50)$$

Because $\vec{p}^2$ is positive-definite, the condition (49) requires for finite κ that $p_0(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = -p'_0(\vec{\mathcal{P}}, \pi'; \vec{\mathcal{Q}}, \rho')$, $q_0(\vec{\mathcal{P}}, \pi; \vec{\mathcal{Q}}, \rho) = -q'_0(\vec{\mathcal{P}}, \pi'; \vec{\mathcal{Q}}, \rho')$ what contradicts the relations (43).

In conclusion, both relations (41) and (43) required for the validity of the relation (34) can not be valid, and consequently if $x_0 \neq y_0$ for on-shell oscillators, it is not possible to factorize in the κ -deformed commutator function (25) any κ -deformed oscillator algebra.

Examples in recent literature. In recent papers [3]-[5] the following particular choice has been made

$$\vec{p} = \vec{\mathcal{P}} \quad (f = 1, g = 0) \quad , \quad \vec{q} = \vec{\mathcal{Q}} \quad (l = 1, k = 0) \quad , \quad p_0 = \mathcal{P}_0 \quad , \quad q_0 = \mathcal{Q}_0. \quad (51)$$

The restrictions on functions $\vec{p}'(\mathcal{P}, \mathcal{Q})$, $\vec{q}'(\mathcal{P}, \mathcal{Q})$ were obtained from non-Abelian composition law of threemomenta (see (24)), considered as an identity in the variables $\vec{\mathcal{P}}, \vec{\mathcal{Q}}$

$$\vec{\mathcal{P}} \dot{+} \vec{\mathcal{Q}} = \vec{q}'(\mathcal{P}, \mathcal{Q}) \dot{+} \vec{p}'(\mathcal{P}, \mathcal{Q}), \quad (52)$$

where $\dot{+}$ denotes addition law based on κ -deformed fourmomentum coproduct. Second relation follows from the energy conservation law.

For our choice of the threemomenta coproduct (see (12)) eq. (52) takes the following explicit form

$$\vec{\mathcal{P}} e^{\frac{\mathcal{P}_0}{2\kappa}} + \vec{\mathcal{Q}} e^{-\frac{\mathcal{Q}_0}{2\kappa}} = \vec{q}'(\mathcal{P}, \mathcal{Q}) e^{\frac{p'_0}{2\kappa}} + \vec{p}'(\mathcal{P}, \mathcal{Q}) e^{-\frac{q'_0}{2\kappa}}, \quad (53)$$

where $\mathcal{P}_0 = \pm\omega_\kappa(\vec{\mathcal{P}})$, $\mathcal{Q}_0 = \pm\omega_\kappa(\vec{\mathcal{Q}})$ and the relations (45) should be inserted on rhs of (53). One gets the relation (53) as identity in particular if⁹

$$\vec{p}'(\mathcal{P}, \mathcal{Q}) = \vec{\mathcal{P}} \exp\left(\frac{\mathcal{P}_0 + q'_0}{2\kappa}\right) \quad , \quad \vec{q}'(\mathcal{P}, \mathcal{Q}) = \vec{\mathcal{Q}} \exp\left(-\frac{\mathcal{Q}_0 + p'_0}{2\kappa}\right), \quad (54)$$

with the relations (45) taking the following explicite form ($\epsilon' = \pm 1, \eta' = \pm 1$)

$$p'_0 = \epsilon' \omega_\kappa(\vec{\mathcal{P}} e^{\frac{\mathcal{P}_0 + q'_0}{2\kappa}}) \quad , \quad q'_0 = \eta' \omega_\kappa(\vec{\mathcal{Q}} e^{-\frac{\mathcal{Q}_0 + p'_0}{2\kappa}}), \quad (55)$$

determining $p'_0 = p'_0(\vec{\mathcal{P}}, \vec{\mathcal{Q}})$ and $q'_0 = q'_0(\vec{\mathcal{P}}, \vec{\mathcal{Q}})$ as functions of threemomenta $\vec{\mathcal{P}}$ and $\vec{\mathcal{Q}}$. The energy conservation relation takes the form

$$\mathcal{P}_0 + \mathcal{Q}_0 = p'_0 + q'_0 \Leftrightarrow \epsilon \omega_\kappa(\vec{\mathcal{P}}) + \eta \omega_\kappa(\vec{\mathcal{Q}}) = \epsilon' \omega_\kappa(\vec{\mathcal{P}} e^{\frac{\mathcal{P}_0 + q'_0}{2\kappa}}) + \eta' \omega_\kappa(\vec{\mathcal{Q}} e^{-\frac{\mathcal{Q}_0 + p'_0}{2\kappa}}). \quad (56)$$

It should be stressed however that even if the relations (56) are valid the relations (43) crucial for obtaining the c -number commutator can not satisfied.

⁹In [3]-[5] there is used different bicrossproduct basis of κ -deformed Poincare algebra, what leads to the coproduct $\Delta(\vec{P}') = \vec{P}' \otimes 1 + e^{-\frac{P'_0}{\kappa}} \otimes \vec{P}'$ and suitable modification of our formulae (53),(54).

One can point out that equations (53) do not specify completely the six functions $p'_i(\mathcal{P}, \mathcal{Q})$, $q'_i(\mathcal{P}, \mathcal{Q})$. In [4], [5] it has been additionally assumed that two products of the oscillators $\hat{A}(\vec{\mathcal{P}})\hat{A}(\vec{\mathcal{Q}})$ and multiplied in flipped order $\hat{A}(\vec{q}', q'_0)\hat{A}(\vec{p}', p'_0)$ transform in the same covariant way under the κ -deformed boost generators. It was shown that:

- i) for $D = 2$ κ -deformed system there exist a unique κ -covariant choice of functions $\vec{p}', \vec{q}'$ consistent with relations (56) and $D = 2$ counterpart of (53),
- ii) in $D = 4$ under analogous assumptions the solution is known only in lowest three orders of $\frac{1}{\kappa}$ perturbation expansion.

The results obtained in [4], [5] providing the κ -covariant set of κ -statistics are very interesting, but because they describe the states with the fourmomenta satisfying standard on-shell conditions (38),(39) they are not consistent with the relation (43) which is necessary for obtaining the field commutator as a c -number.

3 Modified κ -deformed star product and free κ -deformed quantum fields

We see from Sect. 2 that the main problem in obtaining c -number field commutators is the difficulty with getting valid the relations (43). We recall that in Sect. 2 we obtained different on-shell values of $p_0(\mathcal{P}, \mathcal{Q})$, $q_0(\mathcal{P}, \mathcal{Q})$ and $p'_0(\mathcal{P}, \mathcal{Q})$, $q'_0(\mathcal{P}, \mathcal{Q})$ following from different forms of the κ -deformed mass-shell conditions (38) and (39). In this Section we shall change the κ -deformed star product (10) in such a way that both modified mass-shell conditions (38) and (39) will become identical. We will provide two pairs of the same modified energy-momentum dispersion relations for $p_0(\mathcal{P}, \mathcal{Q})$, $p'_0(\mathcal{P}, \mathcal{Q})$ and $q_0(\mathcal{P}, \mathcal{Q})$, $q'_0(\mathcal{P}, \mathcal{Q})$, what allows the validity of the relations (43) and c -number commutation function.

For that purpose we shall use modified κ -deformed star product (15). The corresponding field commutator of κ -deformed free fields (2) takes the form

$$\begin{aligned} [\hat{\varphi}(x), \hat{\varphi}(y)]_{\star_\kappa} &= \frac{1}{(2\pi)^3} \int d^4p d^4q \hat{A}(p)\hat{A}(q)e^{ipx} \star_\kappa e^{iqy} \\ &\quad \cdot \delta(C_\kappa^2(\vec{p}e^{q_0/2\kappa}, p_0) - M^2)\delta(C^2(\vec{q}e^{-p_0/2\kappa}, q_0) - M^2) \\ &\quad - \frac{1}{(2\pi)^3} \int d^4p' d^4q' \hat{A}(q')\hat{A}(p')e^{iq'y} \star_\kappa e^{ip'x} \\ &\quad \cdot \delta(C_\kappa^2(q'_0, \vec{q}'e^{p'_0/2\kappa}) - M^2)\delta(C^2(p'_0, \vec{p}'e^{-q'_0/2\kappa}) - M^2). \end{aligned} \quad (57)$$

Introducing the change of momentum variables (26)-(33) one obtains the formula

$$\begin{aligned} [\hat{\varphi}(x), \hat{\varphi}(y)]_{\star_\kappa} &= \\ &= \frac{1}{(2\pi)^3} \int d^4\mathcal{P} d^4\mathcal{Q} J \left(\begin{smallmatrix} p, q \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix} \right) \hat{A}(p_0, \vec{\mathcal{P}}f + \vec{\mathcal{Q}}g)\hat{A}(q_0, \vec{\mathcal{P}}k + \vec{\mathcal{Q}}l) \\ &\quad \cdot \exp[i(p_0x^0 + q_0y^0)] \exp[-i[(\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g)e^{q_0/2\kappa}\vec{x} + (\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l)e^{-p_0/2\kappa}\vec{y}]] \\ &\quad \cdot \delta(C_\kappa^2(p_0, [\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g]e^{q_0/2\kappa}) - M^2)\delta(C_\kappa^2(q_0, [\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l]e^{-p_0/2\kappa}) - M^2) \\ &\quad - \frac{1}{(2\pi)^3} \int d^4\mathcal{P} d^4\mathcal{Q} J \left(\begin{smallmatrix} p', q' \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix} \right) \hat{A}(q'_0, \vec{\mathcal{P}}k' + \vec{\mathcal{Q}}l')\hat{A}(p'_0, \vec{\mathcal{P}}f' + \vec{\mathcal{Q}}g') \end{aligned} \quad (58)$$

$$\begin{aligned} & \cdot \exp i(p'_0 x^0 + q'_0 y^0) \exp[-i[(\vec{\mathcal{P}}k' + \vec{\mathcal{Q}}l')e^{p'_0/2\kappa}\vec{y} + (\vec{\mathcal{P}}f' + \vec{\mathcal{Q}}g')e^{-q'_0/2\kappa}\vec{x}]] \\ & \cdot \delta(C_\kappa^2(q'_0, [\vec{\mathcal{P}}k' + \vec{\mathcal{Q}}l']e^{p'_0/2\kappa}) - M^2) \delta(C_\kappa^2(p'_0, [\vec{\mathcal{P}}f' + \vec{\mathcal{Q}}g']e^{-q'_0/2\kappa}) - M^2) , \end{aligned}$$

where $J\left(\begin{smallmatrix} p, q \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix}\right)$, $J\left(\begin{smallmatrix} p', q' \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix}\right)$ describe respectively the Jacobians of transformations (28),(29) and (32),(33). We see that by replacement (13) we matched in two consecutive terms in (57) the asymmetry of the star product of exponentials with the asymmetry of mass-shell deltas. After the substitution of relations (41) (expressing $f', g', k', l', p'_0, q'_0$ by f, g, k, l, p_0, q_0) we see that in (58) the products of two mass-shell deltas in two consecutive terms are becoming the same, and one can proceed to factorize the binary algebraic relations describing the κ -deformed oscillator algebra. After inserting the relations (41) the energy values p_0, p'_0 and q_0, q'_0 will satisfy the same mass-shell conditions and therefore it will be consistent to assume the relations (43).

The relations (43) provide necessary conditions for the factorization of κ -deformed algebra. One gets

$$\begin{aligned} [\hat{\varphi}(x), \hat{\varphi}(y)]_{\star_\kappa} = & \int d^4\mathcal{P} d^4\mathcal{Q} \left[J\left(\begin{smallmatrix} p, q \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix}\right) \hat{A}(p_0, \vec{\mathcal{P}}f + \vec{\mathcal{Q}}g) \hat{A}(q_0, \vec{\mathcal{P}}k + \vec{\mathcal{Q}}l) + \right. \\ & \left. - J\left(\begin{smallmatrix} p', q' \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix}\right) \hat{A}(q_0, (\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l)e^{-p_0/\kappa}) \hat{A}(p_0, (\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g)e^{q_0/\kappa}) \right] \\ & \cdot \exp[i(p_0 x^0 + q_0 y^0)] \exp(-i[(\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g)e^{q_0/2\kappa}\vec{x} + (\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l)e^{-p_0/2\kappa}\vec{y}]) \\ & \cdot \delta(C_\kappa^2(p_0, [\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g]e^{q_0/2\kappa}) - M^2) \delta(C_\kappa^2(q_0, [\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l]e^{-p_0/2\kappa}) - M^2) . \end{aligned} \quad (59)$$

Under the integral (59) $\mathcal{P}_0 \equiv \tilde{\pi}(\vec{\mathcal{P}}, \vec{\mathcal{Q}})$ and $\mathcal{Q}_0 \equiv \tilde{\rho}(\vec{\mathcal{P}}, \vec{\mathcal{Q}})$ describe respectively the solutions of the following coupled pair of modified κ -deformed mass-shell conditions

$$C_\kappa^2(p_0, [\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g]e^{q_0/2\kappa}) - M^2 = 0 \quad , \quad C_\kappa^2(q_0, [\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l]e^{-p_0/2\kappa}) - M^2 = 0 . \quad (60)$$

In order to obtain the c -number value of the commutator (58) we should postulate the following general κ -deformed oscillator algebra

$$\begin{aligned} & J\left(\begin{smallmatrix} p, q \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix}\right) \hat{A}(p_0, \vec{\mathcal{P}}f + \vec{\mathcal{Q}}g) \hat{A}(q_0, \vec{\mathcal{P}}k + \vec{\mathcal{Q}}l) \\ & - J\left(\begin{smallmatrix} p', q' \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix}\right) \hat{A}(q_0, (\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l)e^{-p_0/\kappa}) \hat{A}(p_0, (\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g)e^{q_0/\kappa}) = c - \text{number} , \end{aligned} \quad (61)$$

where the functions $p_0 = p_0(\tilde{\pi}, \vec{\mathcal{P}}; \tilde{\rho}, \vec{\mathcal{Q}})$, $q_0 = q_0(\tilde{\pi}, \vec{\mathcal{P}}; \tilde{\rho}, \vec{\mathcal{Q}})$ do satisfy the mass-shell equations (60). We add that the c -number on rhs of (61) should be proportional to the Planck constant $\hbar$.

For classical κ -deformed fields the c -number on rhs of the relation (61) vanishes and in such a case the relations (61) describe the κ -deformed braided oscillators. Substituting in (59) the relation (61) with vanishing c -number one obtains the commutator of κ -braided free fields, describing κ -deformation of classical fields.

$$[\hat{\varphi}^{\text{cl}}(x), \hat{\varphi}^{\text{cl}}(y)]_{\star_\kappa} = 0 . \quad (62)$$

In quantum κ -deformed field theory the nonvanishing c -number on rhs of eq. (61) is proportional to Dirac delta with the argument determined by κ -deformed three-momentum addition law. Such a term in general case is specified in Appendix (see also (67)).

The oscillators $\hat{A}(p_0, \vec{p})$ which are present in the relation (61) carry the fourmomentum $p_\mu = (p_0, \vec{p})$

$$P_\mu \triangleright \hat{A}(p_0, \vec{p}) = p_\mu \hat{A}(p_0, \vec{p}) , \quad (63)$$

restricted by the modified κ -deformed mass-shell conditions (60). The product of two oscillators carry respectively the fourmomenta determined by the coproduct rule (12)

$$P_i \triangleright \left(\hat{A}(p_0, \vec{p}) \hat{A}(q_0, \vec{q}) \right) = \left(p_i e^{\frac{q_0}{2\kappa}} + q_i e^{-\frac{p_0}{2\kappa}} \right) \left(\hat{A}(p_0, \vec{p}) \hat{A}(q_0, \vec{q}) \right) , \quad (64)$$

$$P_0 \triangleright \left(\hat{A}(p_0, \vec{p}) \hat{A}(q_0, \vec{q}) \right) = (p_0 + q_0) \left(\hat{A}(p_0, \vec{p}) \hat{A}(q_0, \vec{q}) \right) . \quad (65)$$

Applying the rules (64),(65) to both products of oscillators occurring in (61) we should obtain the same eigenvalues. We get the following relations:

i) the class of addition laws for threemomenta

$$\vec{p}^{(1+2)} = \left(\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g \right) e^{q_0/2\kappa} + \left(\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l \right) e^{-p_0/2\kappa} = \vec{p}^{(2+1)} , \quad (66)$$

satisfied as identity for arbitrary values of $\vec{p}$ and $\vec{q}$. Such additional law imply the following change of the three-momentum Dirac delta in the oscillator algebra (see also (113),(114) in Appendix)

$$\delta^{(3)}(\vec{\mathcal{P}} - \vec{\mathcal{Q}}) \rightarrow \delta^{(3)} \left(\left[\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g \right] e^{q_0/2\kappa} - \left[\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l \right] e^{-p_0/2\kappa} \right) , \quad (67)$$

ii) the standard addition law for energy

$$p_0^{(1+2)} = p_0 + q_0 = p_0^{(2+1)} . \quad (68)$$

The relations (61) and (66) depend on four functions f, g, k, l ; the values of p_0 and q_0 are determined from the pair of equations (60). Different choices of κ -deformed oscillator algebras (61) can be classified by the corresponding explicite form of threemomentum addition laws described by (66). We add that the relation (61) decomposes into four sets of bilinear relations for κ -deformed creation and annihilation operators (see Appendix).

In order to obtain explicit formulas we shall consider now the particular cases of general relation (61). Firstly we recall the algebraic scheme providing the Abelian addition law [1], [2] and further, we present the framework providing the c -number commutator in the case of the non-Abelian addition law (24) (see also [3]-[5]).

i) *Abelian addition law* [1], [2], [25].

In such a case the functions occurring in (61) are the following

$$f = e^{-\frac{Q_0}{2\kappa}} , \quad g = 0 , \quad k = 0 , \quad l = e^{\frac{P_0}{2\kappa}} ; \quad p_0 = \mathcal{P}_0 , \quad q_0 = \mathcal{Q}_0 . \quad (69)$$

The mass-shell conditions (60) take the form

$$C_\kappa^2(\mathcal{P}_0, \vec{\mathcal{P}}) - M^2 = 0 , \quad C_\kappa^2(\mathcal{Q}_0, \vec{\mathcal{Q}}) - M^2 = 0 , \quad (70)$$

i.e. one should put the following on-shell energy values

$$\mathcal{P}_0^{(\pm)} = \pm \omega_\kappa(\vec{\mathcal{P}}) \quad , \quad \mathcal{Q}_0^{(\pm)} = \pm \omega_\kappa(\vec{\mathcal{Q}}) . \quad (71)$$

One can check that from (66) and (68) we get the Abelian addition laws for threemomenta and energy

$$\vec{\mathcal{P}}^{(1+2)} = \vec{\mathcal{P}} + \vec{\mathcal{Q}} = \vec{\mathcal{P}}^{(2+1)} \quad , \quad \mathcal{P}_0^{(1+2)} = \mathcal{P}_0 + \mathcal{Q}_0 = \mathcal{P}_0^{(2+1)} , \quad (72)$$

where $\mathcal{P}_0, \mathcal{Q}_0$ lie on the mass-shells (70). We point out here that in [1] we have chosen the Abelian addition law (72) as the selection principle for the choice of κ -deformed statistics.

It can be shown that for the choices given by (69) the κ -deformed oscillator algebra (61) can be written in the following standard classical form ($[A, B]_\circ := A \circ B - B \circ A$)¹⁰

$$[a_\kappa(\mathcal{P}), a_\kappa(\mathcal{Q})]_\circ = [a_\kappa^\dagger(\mathcal{P}), a_\kappa^\dagger(\mathcal{Q})]_\circ = 0 \quad , \quad [a_\kappa^\dagger(\mathcal{P}), a_\kappa(\mathcal{Q})]_\circ = 2\Omega_\kappa(\vec{\mathcal{P}})\delta^{(3)}(\vec{\mathcal{P}} - \vec{\mathcal{Q}}) , \quad (73)$$

where the creation and annihilation operators occurring in (73) are ($\mathcal{P}_0^{(\pm)} = \pm \omega_\kappa(\vec{\mathcal{P}})$; $\mathcal{P}_0^+ > 0$)

$$a_\kappa(\mathcal{P}) := \hat{A}(\mathcal{P}_0^{(+)}, \vec{\mathcal{P}}) , \quad a_\kappa^\dagger(\mathcal{P}) := \hat{A}(\mathcal{P}_0^{(-)}, \vec{\mathcal{P}}) = \hat{A}(-\mathcal{P}_0^{(+)}, -\vec{\mathcal{P}}) ; \quad \Omega_\kappa(\vec{\mathcal{P}}) \equiv \kappa \sinh \left(\frac{\omega_\kappa(\vec{\mathcal{P}})}{\kappa} \right) \quad (74)$$

and the κ -deformed $\circ$ -multiplication of two oscillators was given in [1], [2].

The κ -deformed multiplication $\circ$ is defined in such a way that the following relation is valid [25]

$$\hat{\varphi}(x) \hat{\star}_\kappa \hat{\varphi}(y) = \hat{\varphi}(x) \circ \hat{\varphi}(y) , \quad (75)$$

where we define

$$\begin{aligned} \hat{\varphi}(x) \circ \hat{\varphi}(y) = & \frac{1}{(2\pi)^3} \int d^4p d^4q e^{i(p_\mu x^\mu + q_\mu y^\mu)} \hat{A}(p_0, \vec{p}) \circ \hat{A}(q_0, \vec{q}) \cdot \\ & \cdot \delta(C_\kappa^2(p_0, \vec{p}) - M^2) \delta(C_\kappa^2(q_0, \vec{q}) - M^2) . \end{aligned} \quad (76)$$

Subsequently

$$[\hat{\varphi}(x), \hat{\varphi}(x)]_{\star_\kappa} = [\hat{\varphi}(x), \hat{\varphi}(y)]_\circ , \quad (77)$$

what after insertion of the relations (73) permits to obtain the c -number value of field commutator described by the formulae (21), (22).

One can point out that in consistency with the classical fourmomenta addition law (72) the choice (69) is the one which provides the particular c -number field commutator function (22) which is invariant under the classical fourdimensional translations $x_\mu \rightarrow x_\mu + a_\mu$, i.e. we obtain the κ -deformed commutator function depending on the four-coordinate difference $x_\mu - y_\mu$ [12].

ii) *Non-Abelian κ -deformed addition law and κ -deformed flip operator.*

We choose now the arbitrary functions in (61) as follows

$$f = 1 \quad , \quad g = 0 \quad , \quad k = 0 \quad , \quad l = 1 \quad ; \quad p_0 = \mathcal{P}_0 \quad , \quad q_0 = \mathcal{Q}_0 . \quad (78)$$

¹⁰In fact the rhs of second relation (73) is multiplied by Planck constant $\hbar$. In our consideration we put $\hbar = 1$.

The mass-shell conditions (60) take the form

$$C_\kappa^2(\mathcal{P}_0, \vec{\mathcal{P}} e^{\mathcal{Q}_0/2\kappa}) - M^2 = 0 \quad , \quad C_\kappa^2(\mathcal{Q}_0, \vec{\mathcal{Q}} e^{-\mathcal{P}_0/2\kappa}) - M^2 = 0 \quad , \quad (79)$$

or more explicitly

$$\mathcal{P}_0 = \epsilon \omega_\kappa(\vec{\mathcal{P}} e^{\mathcal{Q}_0/2\kappa}) \quad , \quad \mathcal{Q}_0 = \epsilon' \omega_\kappa(\vec{\mathcal{Q}} e^{-\mathcal{P}_0/2\kappa}) \quad . \quad (80)$$

Then one gets from (61) the following uniquely determined κ -deformed oscillator algebra

$$\hat{A}(\mathcal{P}_0, \vec{\mathcal{P}}) \hat{A}(\mathcal{Q}_0, \vec{\mathcal{Q}}) - J \left(\vec{\mathcal{P}}, \vec{\mathcal{Q}} \right)^{p', q'} \hat{A}(\mathcal{Q}_0, \vec{\mathcal{Q}} e^{-\frac{\mathcal{P}_0}{\kappa}}) \hat{A}(\mathcal{P}_0, \vec{\mathcal{P}} e^{\frac{\mathcal{Q}_0}{\kappa}}) = c - \text{number} \quad , \quad (81)$$

where the c -number in (81) is proportional to the following Dirac delta

$$\delta^{(3)}(\vec{\mathcal{P}} - \vec{\mathcal{Q}}) \rightarrow \delta^{(3)}(\vec{\mathcal{P}} e^{\mathcal{Q}_0/2\kappa} - \vec{\mathcal{Q}} e^{-\mathcal{P}_0/2\kappa}) \quad , \quad (82)$$

and occurs only in creation-annihilation (or annihilation-creation) sector (see Appendix, (113), (114)). The formula (81) can be written as well with the use of the κ -deformed flip operator

$$\hat{A}(\mathcal{P}_0, \vec{\mathcal{P}}) \hat{A}(\mathcal{Q}_0, \vec{\mathcal{Q}}) - \hat{\tau}_\kappa \left(\hat{A}(\mathcal{P}_0, \vec{\mathcal{P}}) \hat{A}(\mathcal{Q}_0, \vec{\mathcal{Q}}) \right) = c - \text{number} \quad , \quad (83)$$

where

$$\hat{\tau}_\kappa \left(\hat{A}(\mathcal{P}_0, \vec{\mathcal{P}}) \hat{A}(\mathcal{Q}_0, \vec{\mathcal{Q}}) \right) = J \left(\vec{\mathcal{P}}, \vec{\mathcal{Q}} \right)^{p', q'} \hat{A}(\mathcal{Q}_0, \vec{\mathcal{Q}} e^{-\frac{\mathcal{P}_0}{\kappa}}) \hat{A}(\mathcal{P}_0, \vec{\mathcal{P}} e^{\frac{\mathcal{Q}_0}{\kappa}}) \quad , \quad (84)$$

and $\mathcal{P}_0, \mathcal{Q}_0$ are the solutions of the equations (80). Using twice the formula (84) and the property $J \left(\vec{\mathcal{P}}, \vec{\mathcal{Q}} \right)^{p, q} J \left(\vec{\mathcal{P}}, \vec{\mathcal{Q}} \right) = 1$ it is easy to see that the flip operator $\hat{\tau}_\kappa$ satisfies the condition

$$\hat{\tau}_\kappa^2 = 1 \quad . \quad (85)$$

In order to adjust the set of relations (111)-(114) (see Appendix) to the choice (78) of functions f, g, k, l, p_0 and q_0 we should solve the equations (80). The equation for the solutions $\mathcal{P}_0^{(\epsilon, \epsilon')} = \mathcal{P}_0^{(\epsilon, \epsilon')}(\vec{\mathcal{P}}, \vec{\mathcal{Q}})$ looks as follows

$$\mathcal{P}_0^{(\epsilon, \epsilon')} = \epsilon \omega_\kappa \left(\vec{\mathcal{P}} \exp \left[\frac{\epsilon'}{2\kappa} \omega_\kappa \left(\vec{\mathcal{Q}} \exp(-\mathcal{P}_0^{(\epsilon, \epsilon')}/2\kappa) \right) \right] \right) \quad . \quad (86)$$

One gets the second set of energy values using the relation

$$\mathcal{Q}_0^{(\epsilon, \epsilon')}(\vec{\mathcal{Q}}, \vec{\mathcal{P}}; \kappa) = \mathcal{P}_0^{(\epsilon, \epsilon')}(\vec{\mathcal{P}}, \vec{\mathcal{Q}}; -\kappa) \quad . \quad (87)$$

The formulae for $\mathcal{P}_0^{(\epsilon, \epsilon')}$ and $\mathcal{Q}_0^{(\epsilon, \epsilon')}$ after inserting in (81) provide explicit example of four set (111)-(114) (see Appendix) of the κ -deformed twisted oscillator algebra for creation and annihilation operators. After inserting these algebraic relations in field commutator (59) we can show that the commutator (57) one obtains a c -number value.

4 General algebraic structure of κ -deformed oscillator algebras

Let us consider firstly standard undeformed theory. The standard algebra describing the bosonic oscillators $a(\mathcal{P}), a^\dagger(\mathcal{Q})$ looks as follows

$$[a^\dagger(\mathcal{P}), a^\dagger(\mathcal{Q})] = [a^\dagger(\mathcal{P}), a^\dagger(\mathcal{Q})] = 0 \quad , \quad [a^\dagger(\mathcal{P}), a^\dagger(\mathcal{Q})] = 2\omega(\vec{\mathcal{P}})\delta^3(\vec{\mathcal{P}} - \vec{\mathcal{Q}}), \quad (88)$$

where $\mathcal{P} \equiv (\mathcal{P}_0 = \omega(\vec{\mathcal{P}}), \vec{\mathcal{P}})$.

The κ -deformation of the algebra (88) can be introduced in two ways:

- i) *The deformation of oscillator algebra (88) described by the general κ -deformed multiplication rule.*

In order to describe the general algebra (61) we introduce new κ -deformed multiplication $\odot$ generalizing the $\circ$ -multiplication given in [1], [2] as follows

$$\begin{aligned} \hat{A}(\mathcal{P})|_{\mathcal{P}_0=\epsilon\omega_\kappa(\vec{\mathcal{P}})} \odot \hat{A}(\mathcal{Q})|_{\mathcal{P}_0=\epsilon'\omega_\kappa(\vec{\mathcal{P}})} &:= J \left(\begin{matrix} p(\mathcal{P}, \mathcal{Q}), q(\mathcal{P}, \mathcal{Q}) \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \\ &\cdot \hat{A}(p_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}), \vec{p}(\mathcal{P}, \mathcal{Q})) \hat{A}(q_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}), \vec{q}(\mathcal{P}, \mathcal{Q}))|_{\mathcal{P}_0=\tilde{\pi}, \mathcal{Q}_0=\tilde{\rho}}, \end{aligned} \quad (89)$$

where $\epsilon = \epsilon' = 1$ describes the creation-creation sector, $\epsilon = -\epsilon'$ the creation-annihilation sector, and $\epsilon = \epsilon' = -1$ the annihilation-annihilation sector.

We described in such a way the first part of binary relation (61) depending on six arbitrary functions. One can consider a subclass of the multiplication rule (89) which permit to describe the relation (61) as the standard oscillator algebra (88) with the generalized κ -deformed multiplication. For such a purpose the multiplication (89) should satisfy additional relation permitting as well to express second part of the binary relation (61) by the use of multiplication rule (89), namely

$$\begin{aligned} \hat{A}(\mathcal{Q})|_{\mathcal{Q}_0=\epsilon'\omega_\kappa(\vec{\mathcal{Q}})} \odot \hat{A}(\mathcal{P})|_{\mathcal{P}_0=\epsilon\omega_\kappa(\vec{\mathcal{P}})} &= J \left(\begin{matrix} p'(\mathcal{P}, \mathcal{Q}), q'(\mathcal{P}, \mathcal{Q}) \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \\ &\cdot \hat{A}(q_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}), \vec{q}(\mathcal{P}, \mathcal{Q})) e^{-p_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q})/\kappa} \\ &\cdot \hat{A}(p_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}), \vec{p}(\mathcal{P}, \mathcal{Q})) e^{q_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q})/\kappa} |_{\mathcal{P}_0=\tilde{\pi}, \mathcal{Q}_0=\tilde{\rho}}. \end{aligned} \quad (90)$$

The validity of (90) restricts six arbitrary functions occurring in (61) in the following way

$$p_i^T(\mathcal{P}, \mathcal{Q}) \equiv p_i(\mathcal{Q}, \mathcal{P}) = q_i(\mathcal{P}, \mathcal{Q}) e^{-p_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q})/\kappa} \quad , \quad (p_0^{(\epsilon, \epsilon')})^T(\mathcal{P}, \mathcal{Q}) = q_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}), \quad (91)$$

$$q_i^T(\mathcal{P}, \mathcal{Q}) \equiv q_i(\mathcal{Q}, \mathcal{P}) = p_i(\mathcal{P}, \mathcal{Q}) e^{q_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q})/\kappa} \quad , \quad (q_0^{(\epsilon, \epsilon')})^T(\mathcal{P}, \mathcal{Q}) = p_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}). \quad (92)$$

The algebra (61) with the choice (91), (92) can be written as follows

$$[a(\mathcal{P}), a(\mathcal{Q})]_\odot = [a^\dagger(\mathcal{P}), a^\dagger(\mathcal{Q})]_\odot = 0 \quad , \quad [a^\dagger(\mathcal{P}), a(\mathcal{Q})]_\odot = c - \text{number}. \quad (93)$$

The relations (93) describe the class of κ -deformed oscillator algebras with generalized κ -deformed multiplication, parametrized by three arbitrary functions solving the conditions (91),(92)

In particular if $p_0 = \mathcal{P}_0, q_0 = \mathcal{Q}_0$ and we choose (see (69))

$$\vec{p}(\mathcal{P}, \mathcal{Q}) = e^{-\frac{\mathcal{Q}_0}{2\kappa}} \vec{\mathcal{P}} \quad , \quad \vec{q}(\mathcal{P}, \mathcal{Q}) = e^{\frac{\mathcal{P}_0}{2\kappa}} \vec{\mathcal{Q}} \quad , \quad (94)$$

we get

$$\vec{p}^T(\mathcal{P}, \mathcal{Q}) = e^{-\frac{\mathcal{P}_0}{2\kappa}} \vec{\mathcal{Q}} \quad , \quad \vec{q}^T(\mathcal{P}, \mathcal{Q}) = e^{\frac{\mathcal{Q}_0}{2\kappa}} \vec{\mathcal{P}} \quad , \quad (95)$$

where $\mathcal{P}_0 = \mathcal{P}_0^{(\epsilon)}$, $\mathcal{Q}_0 = \mathcal{Q}_0^{(\epsilon')}$.

We see that the choice (69) which provides first example of κ -multiplication [1] obviously satisfies the conditions (91),(92).

ii) *The general κ -deformed flip operator.*

Let us describe the commutator of two standard bosonic oscillators as follows

$$[\hat{A}(\mathcal{P}), \hat{A}(\mathcal{Q})] \equiv \hat{A}(\mathcal{P})\hat{A}(\mathcal{Q}) - \hat{\tau}_0[\hat{A}(\mathcal{P})\hat{A}(\mathcal{Q})] = c - \text{number} \quad , \quad (96)$$

where $\mathcal{P}_0 = \epsilon\omega(\vec{\mathcal{P}})$ and $\hat{\tau}_0(\hat{A}\hat{B}) = \hat{B}\hat{A}$. In particular if $n = 2$, the symmetrization operator S_2 is described in terms of flip operator as follows

$$S_2^{(0)} = \frac{1}{2}(1 + \hat{\tau}_0) \quad . \quad (97)$$

One can look for the κ -deformation of the classical flip operator $\hat{\tau}_0$, which leads to the following κ -deformation of the commutator (96)

$$[\hat{A}, \hat{B}] \rightarrow [\hat{A}, \hat{B}]_{\hat{\tau}_\kappa} := \hat{A}\hat{B} - \hat{\tau}_\kappa\hat{A}\hat{B} \quad , \quad (98)$$

where we should assume that

$$[\hat{\tau}_\kappa, \Delta(P_\mu)] = 0 \quad . \quad (99)$$

We supplement the multiplication (89) with the following definition of κ -deformed flip operation

$$\begin{aligned} & \hat{\tau}_\kappa \left[\hat{A}(\mathcal{P})|_{\mathcal{P}_0=\epsilon\omega_\kappa(\vec{\mathcal{P}})} \odot \hat{A}(\mathcal{Q})|_{\mathcal{Q}_0=\epsilon'\omega_\kappa(\vec{\mathcal{Q}})} \right] \\ & \equiv \hat{\tau}_\kappa \left[J \left(\begin{matrix} p(\mathcal{P}, \mathcal{Q}), q(\mathcal{P}, \mathcal{Q}) \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \hat{A}(p_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}), \vec{p}(\mathcal{P}, \mathcal{Q})) \hat{A}(q_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}), \vec{q}(\mathcal{P}, \mathcal{Q})) \right] |_{\mathcal{P}_0=\tilde{\pi}, \mathcal{Q}_0=\tilde{\rho}} \\ & := J \left(\begin{matrix} p'(\mathcal{P}, \mathcal{Q}), q'(\mathcal{P}, \mathcal{Q}) \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \hat{A}(q_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}), \vec{q}(\mathcal{P}, \mathcal{Q})) e^{-p_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q})/\kappa} \\ & \quad \cdot \hat{A}(p_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q}), \vec{p}(\mathcal{P}, \mathcal{Q})) e^{q_0^{(\epsilon, \epsilon')}(\mathcal{P}, \mathcal{Q})/\kappa} |_{\mathcal{P}_0=\tilde{\pi}, \mathcal{Q}_0=\tilde{\rho}} \quad , \end{aligned} \quad (100)$$

consistently with (99). Further, one can show that for any choice of the multiplication $\odot$ we obtain

$$\hat{\tau}_\kappa^2 = 1 \quad . \quad (101)$$

In particular one can consider a κ -statistics described only by κ -deformed flip operator with standard undeformed multiplication rule

$$\hat{A}(\mathcal{P}) \odot \hat{A}(\mathcal{Q}) \equiv \hat{A}(\mathcal{P})\hat{A}(\mathcal{Q}). \quad (102)$$

Such a case is obtained by putting $f = l = 1$, $g = k = 0$ and $p_0 = \mathcal{P}_0$, $q_0 = \mathcal{Q}_0$, and the flip operator is uniquely defined as follows (see (84))

$$\hat{\tau}_\kappa[\hat{A}(\mathcal{P})\hat{A}(\mathcal{Q})] = J\left(\begin{smallmatrix} p' & q' \\ \vec{\mathcal{P}} & \vec{\mathcal{Q}} \end{smallmatrix}\right) \hat{A}(\mathcal{Q}_0, \vec{\mathcal{Q}}e^{-\frac{\mathcal{P}_0}{\kappa}})\hat{A}(\mathcal{P}_0, \vec{\mathcal{P}}e^{\frac{\mathcal{Q}_0}{\kappa}}). \quad (103)$$

iii) *The most general κ -deformed oscillator algebra.*

We shall consider now the most general κ -deformed oscillator algebra (61), which for the energy-momentum dispersion relation described by the mass-shell conditions (60) leads to c -number commutator function. It appears that such algebra can be described by the composition of the multiplication (89) and κ -deformed flip (100) in the following way

$$(1 - \hat{\tau}_\kappa) \left[\hat{A}(\mathcal{P})|_{\mathcal{P}_0=\epsilon\omega_\kappa(\vec{\mathcal{P}})} \odot \hat{A}(\mathcal{Q})|_{\mathcal{Q}_0=\epsilon'\omega_\kappa(\vec{\mathcal{Q}})} \right] = c - \text{number}. \quad (104)$$

If we describe the 2-particle sector of standard bosonic Hilbert space $\mathcal{H}_2$ as symmetrized tensor product of one particle Hilbert spaces $\mathcal{H}_1$

$$\mathcal{H}_2 = S_2 \circ (\mathcal{H}_1 \otimes \mathcal{H}_1), \quad (105)$$

the κ -deformed multiplication modifies the tensor product as follows

$$\mathcal{H}_1 \otimes \mathcal{H}_1 \rightarrow \mathcal{H}_1 \otimes_\kappa \mathcal{H}_1, \quad (106)$$

where $\otimes_\kappa$ is obtained from the "braided multiplication" $\odot$ of oscillators (see (89)) and κ -deformed twist $\hat{\tau}_\kappa$ changes the symmetrization operator

$$S_2 \rightarrow S_2^\kappa = \frac{1}{2}(1 + \hat{\tau}_\kappa). \quad (107)$$

In this paper we limited our considerations to the most general binary algebraic relations. Only in the case of algebra (61) with the particular choice of arbitrary functions given by (69) (see [1], [2]) the products of arbitrary number of κ -deformed creation and annihilation operators have been introduced, and all sectors of corresponding κ -deformed Hilbert space have been considered.

5 Final Remarks

In this paper we consider the general structure of κ -deformed binary oscillator algebras and their applicability for description of free κ -deformed quantum fields, with c -number commutator functions. Our considerations generalize recent examples of κ -deformed statistics studied

by present authors [1], [2], [25], Arzano and Marciano [3], Young and Zegers [4], [5] and Govindarajan et al [16]. We consider two different classes of the κ -deformed oscillators which differ by their κ -deformed energy-momentum relations: in Sect. 2 we assume that the oscillators $\hat{A}(p_0, \vec{p})$ are put on standard κ -deformed mass shell (4), and in Sect. 3 we modify the standard κ -deformed mass-shell conditions in a way providing c -number field commutators.

The basic results obtained in this paper are the following:

- i) We considered in Sect. 2 the algebra of deformed oscillators on κ -deformed mass-shell, corresponding to noncommutative field theory with standard multiplication of noncommutative fields, via standard κ -star product (10). In such a case for arbitrary choice of κ -deformed oscillator algebra the κ -deformed quantum fields are not free. One can show that the commutator function is necessarily bilinear in field operators.
- ii) We studied in Sect. 3 the general algebra of oscillators with modified κ -deformed mass-shell conditions (see (79)), corresponding to noncommutative field theory with modified multiplication law of noncommutative fields (see (14)) represented by nonstandard κ -star product (15). In such a case one obtains for large class of κ -deformed oscillator algebras (see (61) and Appendix) the κ -deformed free quantum fields which are characterized by the c -number commutator function. It should be stressed that only in such a case one can look for the formulation of the κ -deformed perturbative description of interacting κ -deformed field theory and consider the suitable generalization of Feynman diagram technique.
- iii) In Sect. 4 we did show that the general κ -deformed statistics providing κ -deformed quantum fields with c -number commutators is obtained by the composition of general κ -deformed multiplication and the κ -deformed flip operator.

We point out finally that in this paper we do not discuss the κ -covariance of the κ -deformed oscillator algebras. The results of [4], [5] suggest however that the κ -covariance is not consistent with the relation (43) which we had to postulate in order to obtain the c -number κ -fields commutator. It appears therefore that the freedom of choice present in the formula (61) might not permit to obtain both the manifest κ -covariance as well as the c -number commutator function.

6 Appendix: General algebra of κ -deformed creation and annihilation operators

One can rewrite the mass-shell conditions (60) as the set of four nonlinear algebraic equations describing eight classes of energy-momentum dispersion relations $p_0 \equiv p_0^{(\epsilon, \epsilon')}(\tilde{\pi}, \vec{\mathcal{P}}; \tilde{\rho}, \vec{\mathcal{Q}})$, $q_0 \equiv q_0^{(\epsilon, \epsilon')}(\tilde{\pi}, \vec{\mathcal{P}}; \tilde{\rho}, \vec{\mathcal{Q}})$ ($\epsilon = \pm 1$, $\epsilon' = \pm 1$)

$$p_0 = \epsilon \omega_\kappa([\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g]e^{q_0/2\kappa}) \quad , \quad q_0 = \epsilon' \omega_\kappa([\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l]e^{-p_0/2\kappa}) \quad , \quad (108)$$

where

$$\lim_{\kappa \rightarrow \infty} p_0^{(\epsilon, \epsilon')} = \epsilon \omega(\vec{p}) \quad , \quad \lim_{\kappa \rightarrow \infty} q_0^{(\epsilon, \epsilon')} = \epsilon' \omega(\vec{q}) \quad . \quad (109)$$

The values $p_0^{(1, \epsilon')} > 0$, $(q_0^{(\epsilon, 1)} > 0)$ describe the on-shell values of p_0 (q_0) which select $\hat{A}(p_0^{(1, \epsilon')}, \vec{p})$, $\hat{A}(q_0^{(\epsilon, 1)}, \vec{q})$ as κ -deformed creation operators; the on-shell energy values $p_0^{(-1, \epsilon')} < 0$, $(q_0^{(\epsilon, -1)} < 0)$ are required to define the κ -deformed annihilation operators $\hat{A}(p_0^{(-1, \epsilon')}, \vec{p})$, $\hat{A}(q_0^{(\epsilon, -1)}, \vec{q})$. If we denote

$$p^{(\epsilon, \epsilon')} = (p_0^{(\epsilon, \epsilon')}, \vec{p}(\tilde{\pi}, \vec{P}; \tilde{\rho}, \vec{Q})) \quad , \quad q^{(\epsilon, \epsilon')} = (q_0^{(\epsilon, \epsilon')}, \vec{q}(\tilde{\pi}, \vec{P}; \tilde{\rho}, \vec{Q})) \quad , \quad (110)$$

one can rewrite the relations (61) as the four set of relations describing creation-creation, creation-annihilation, annihilation-creation and annihilation-annihilation sectors.

If we insert in proper way the on-shell energy values $p_0^{(\epsilon, \epsilon')}$, $q_0^{(\epsilon, \epsilon')}$ (see (108),(110)) the relations (61) decompose into the following set of relations:

i) creation-creation algebra

$$\begin{aligned} & J \left(\begin{matrix} p^{(++)}, q^{(++)} \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \hat{A}(p_0^{(++)}, \vec{P}f + \vec{Q}g) \cdot \hat{A}(q_0^{(++)} \vec{P}k + \vec{Q}l) \\ & = J \left(\begin{matrix} p'^{(++)}, q'^{(++)} \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \hat{A}(q_0^{(++)}, [\vec{P}k + \vec{Q}l]e^{-p_0^{(++)}/\kappa}) \cdot \hat{A}(p_0^{(++)}, [\vec{P}f + \vec{Q}g]e^{q_0^{(++)}/\kappa}) \quad , \end{aligned} \quad (111)$$

ii) annihilation-annihilation algebra

$$\begin{aligned} & J \left(\begin{matrix} p^{(--)}, q^{(--)} \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \hat{A}(p_0^{(--)}, \vec{P}f + \vec{Q}g) \cdot \hat{A}(q_0^{(--)} \vec{P}k + \vec{Q}l) \\ & = J \left(\begin{matrix} p'^{(--)}, q'^{(--)} \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \hat{A}(q_0^{(--)}, [\vec{P}k + \vec{Q}l]e^{-p_0^{(--)} / \kappa}) \cdot \hat{A}(p_0^{(--)}, [\vec{P}f + \vec{Q}g]e^{q_0^{(--)} / \kappa}) \quad , \end{aligned} \quad (112)$$

iii) creation-annihilation algebra

$$\begin{aligned} & J \left(\begin{matrix} p^{(+-)}, q^{(-+)} \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \hat{A}(p_0^{(+-)}, \vec{P}f + \vec{Q}g) \cdot \hat{A}(q_0^{(-+)} \vec{P}k + \vec{Q}l) \\ & - J \left(\begin{matrix} p'^{(-+)}, q'^{(-+)} \\ \mathcal{P}, \mathcal{Q} \end{matrix} \right) \hat{A}(q_0^{(-+)}, [\vec{P}k + \vec{Q}l]e^{-p_0^{(-+)} / \kappa}) \cdot \hat{A}(p_0^{(+-)}, [\vec{P}f + \vec{Q}g]e^{q_0^{(-+)} / \kappa}) \\ & = 2N_{\kappa}^{(+ -)}(\vec{P}, \vec{Q}) \delta^{(3)} \left[(\vec{P}f + \vec{Q}g)e^{q_0^{(-+)} / \kappa} - (\vec{P}k + \vec{Q}l)e^{-p_0^{(+ -)} / \kappa} \right] \quad , \end{aligned} \quad (113)$$

iv) annihilation-creation algebra

$$\begin{aligned}
& J \left(\begin{smallmatrix} p^{(-+)} \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix} \right) \hat{A}(p_0^{(-+)}, \vec{\mathcal{P}}f + \vec{\mathcal{Q}}g) \cdot \hat{A}(q_0^{(+-)} \vec{\mathcal{P}}k + \vec{\mathcal{Q}}l) \\
& - J \left(\begin{smallmatrix} p'^{(-+)} \\ \vec{\mathcal{P}}, \vec{\mathcal{Q}} \end{smallmatrix} \right) \hat{A}(q_0^{(+-)}, [\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l]e^{-p_0^{(-+)}/\kappa}) \cdot \hat{A}(p_0^{(-+)}, [\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g]e^{q_0^{(+-)}/\kappa}) \\
& = 2N_\kappa^{(-+)}(\vec{\mathcal{P}}, \vec{\mathcal{Q}}) \delta^{(3)} \left[(\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g)e^{q_0^{(+-)}/\kappa} - (\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l)e^{-p_0^{(-+)}/\kappa} \right] .
\end{aligned} \tag{114}$$

The threemomentum Dirac delta in (113),(114) describes in accordance with the formula (66) the κ -deformed threemomentum conservation law for the process of creation/annihilation of field quanta with the threemomentum $(\vec{\mathcal{P}}f + \vec{\mathcal{Q}}g)$ and annihilation of the quanta with the momentum $(\vec{\mathcal{P}}k + \vec{\mathcal{Q}}l)$.

Acknowledgments

This paper has been financially supported by Polish Ministry of Science and Higher Education grant NN202318534.

References

- [1] M. Daszkiewicz, J. Lukierski, M. Woronowicz, Mod. Phys. Lett. A 23, 9 (2008); hep-th/0703200
- [2] M. Daszkiewicz, J. Lukierski, M. Woronowicz, Phys. Rev. D 77, 105007 (2008); arXiv: 0708.1561 [hep-th]
- [3] M. Arzano, A. Marciano, Phys. Rev. D 76, 125005 (2007); arXiv:0707.1329 [hep-th]
- [4] C.A.S. Young, R. Zegers, Nucl. Phys. B 797, 537 (2008); arXiv: 0711.2206 [hep-th]
- [5] C.A.S. Young, R. Zegers, "Covariant particle exchange for κ -deformed theories in 1+1 dimensions"; arXiv: 0803.2659 [hep-th]
- [6] S. Doplicher, K. Fredenhagen, J.E. Roberts, Phys. Lett. B 331, 39 (1994)
- [7] S. Doplicher, K. Fredenhagen, J.E. Roberts, Comm. Math. Phys. 172, 187 (1995)
- [8] S. Zakrzewski, J. Phys. A 27, 2075 (1994)
- [9] H. Ruegg and S. Majid, Phys. Lett. B 334, 348 (1994)
- [10] J. Lukierski, H. Ruegg, W.J. Zakrzewski, Annals of Phys. 243, 90 (1995)
- [11] J. Lukierski, A. Nowicki, H. Ruegg and V.N. Tolstoy, Phys. Lett. B 264, 331 (1991)

- [12] J. Lukierski, A. Nowicki, H. Ruegg, Phys. Lett. B 293, 344 (1992)
- [13] M. Chaichian, P.P. Kulish, K. Nishijima, A. Turenau, Phys. Lett. B 604, 98 (2004)
- [14] G. Fiore, J. Wess; Phys. Rev D 75, 105022 (2007); hep-th/0701078
- [15] J.G. Bu, H. Ch. Kim, Y. Lee, C.H. Vac, J.H. Yee, Phys. Lett. B 665, 95 (2008); hep-th/0611175
- [16] T.R. Govindarajan, Kumar S. Gupta, E. Harikumar, S. Meljanac and D. Meljanac, Phys. Rev. D 77, 105010 (2008); arXiv: 0802.1576 [hep-th]
- [17] P. Kosinski, J. Lukierski, P. Maslanka, Czech. J. Phys. 50, 1283 (2000)
- [18] G. Amelino-Camelia, M. Arzano, Phys. Rev. D 65, 084044 (2002); hep-th/0105120
- [19] M. Dimitrijevič, L. Jonke, L. Möller, E. Tsouchnika, J. Wess, M. Wohlegennut, Euro. Phys. J C 31, 129 (2003); hep-th/0307149
- [20] M. Dimitrijevič, L. Möller, E. Tsouchnika, J. Phys. A 37, 9749 (2004); hep-th/0404224
- [21] L. Freidel, J. Kowalski-Glikman and S. Nowak, Phys. Lett. B 648, 70 (2007)
- [22] S. Meljanac, A. Sensorov, M. Stojic and Kumar S. Gupta, Eur. Phys. J. C 53, 295 (2008); arXiv: 0705.2471 [hep-th]
- [23] A.L. Licht and J.S. Toll, Nuovo Cim. 21, 346 (1961)
- [24] R. Jost, *"The general theory of quantized fields"*; Providence, Rhode Island, Amer. Math. Soc., 1965
- [25] M. Daszkiewicz, J. Lukierski, M. Woronowicz, *"Quantization of κ -deformed free fields and κ -deformed oscillators"*; Proc. of 7th International Workshop on Supersymmetries and Quantum Symmetries (SQS'07), Dubna, Russia, August 2007, ed. S. Fedoruk and E. Ivanov, ed. JINR Dubna, p. 197 (2008); arXiv: 0712.0350 [hep-th]
- [26] A. Agostini, F. Lizzi, A. Zampini, Mod. Phys. Lett. A 17, 2105 (2002); hep-th/0209174
- [27] D. Bahns, S. Doplicher, K. Fredenhagen, G. Praditielli, Comm. Math. Phys. 237, 221 (2003); hep-th/0301100
- [28] J. Lukierski, *" κ -deformed oscillators: deformed multiplication versus deformed flip operator and multiparticle clusters"*; arXiv: 0812.0547 [hep-th]; Reports in Math. Physics, in press.